


Utah Water Supply Outlook Report

Jan, 2005



Mike Bricco measures snow at Smith and Morehouse SNOTEL site, December 22, 2004. Photo by Randy Julander

Water Supply Outlook Reports and Federal - State - Private Cooperative Snow Surveys

For more water supply and resource management information, contact:

Vane O. Campbell, Area Conservationist, 340 N. 600 E., Richfield, UT 84701 - Phone: (435) 896-6441
Todd C. Nielson, Area Conservationist, 302 E. 1860 S., Provo, UT 84606 - Phone: (801) 377

Todd C. Nielson, Area Conservationist, 302 E. 1860 S., Provo, UT 84606 - Phone: (801) 377-5580 David M. Webster, Area Conservationist, 80 N. 500 W., Vernal, UT 84078 - Phone: (435)789-2100

Snow Survey Staff, 245 N Jimmy Doolittle Rd, SLC Utah, 84041 - Phone: (801)524-5213

Internet Address: http://www.ut.nrcs.usda.gov/snow/

How forecasts are made

Most of the annual streamflow in the western United States originates as snowfall that has accumulated in the mountains during the winter and early spring. As the snowpack accumulates, hydrologists estimate the runoff that will occur when it melts. Measurements of snow water equivalent at selected manual snowcourses and automated SNOTEL sites, along with precipitation, antecedent streamflow, and indices of the El Niño / Southern Oscillation are used in computerized statistical and simulation models to prepare runoff forecasts. These forecasts are coordinated between hydrologists in the Natural Resources Conservation Service and the National Weather Service. Unless otherwise specified, all forecasts are for flows that would occur naturally without any upstream influences.

Forecasts of any kind, of course, are not perfect. Streamflow forecast uncertainty arises from three primary sources: (1) uncertain knowledge of future weather conditions, (2) uncertainty in the forecasting procedure, and (3) errors in the data. The forecast, therefore, must be interpreted not as a single value but rather as a range of values with specific probabilities of occurrence. The middle of the range is expressed by the 50% exceedance probability forecast, for which there is a 50% chance that the actual flow will be above, and a 50% chance that the actual flow will be below, this value. To describe the expected range around this 50% value, four other forecasts are provided, two smaller values (90% and 70% exceedance probability) and two larger values (30%, and 10% exceedance probability). For example, there is a 90% chance that the actual flow will be more than the 90% exceedance probability forecast. The others can be interpreted similarly.

The wider the spread among these values, the more uncertain the forecast. As the season progresses, forecasts become more accurate, primarily because a greater portion of the future weather conditions become known; this is reflected by a narrowing of the range around the 50% exceedance probability forecast. Users should take this uncertainty into consideration when making operational decisions by selecting forecasts corresponding to the level of risk they are willing to assume about the amount of water to be expected. If users anticipate receiving a lesser supply of water, or if they wish to increase their chances of having an adequate supply of water for their operations, they may want to base their decisions on the 90% or 70% exceedance probability forecasts, or something in between. On the other hand, if users are concerned about receiving too much water (for example, threat of flooding), they may want to base their decisions on the 30% or 10% exceedance probability forecasts, or something in between. Regardless of the forecast value users choose for operations, they should be prepared to deal with either more or less water. (Users should remember that even if the 90% exceedance probability forecast is used, there is still a 10% chance of receiving less than this amount.) By using the exceedance probability information, users can easily determine the chances of receiving more or less water.

The United States Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326 W, Whitten Building, 14th and Independence Ave., SW, Washington, D.C., 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

STATE OF UTAH GENERAL OUTLOOK Jan 1, 2005

SUMMARY

Water year 2005 has distinct similarities to the beginning of water year 2004 as well as some distinct differences. Last year had some phenomenal storms that jump started the snowpacks across the state and the same is true for this year. In northern Utah on the Bear, Weber and Provo watersheds, current snowpacks (110% to 140%) are a mirror image of last year and that is pretty much where the similarity ends. The Uintah basin and North Slope snowpacks are at 177% of normal, about 50% more than last year. The Uintah basin currently has 6 sites that have over 200% of average snowpack. This area has a 97% probability of at least average snowpack by April 1 and a significant probability of snowpacks greater than 130%. Further south, the Sevier River (172%) and most of southern Utah (246%) have very impressive snowpacks. The Sevier has an 85% probability of at least average snowpack by April 1 and southwestern Utah has an 88% probability. Given average snowpack accumulation over the remainder of winter, these areas will be between 130 and 160% of average on April 1. Both of these areas have current snowpacks similar to the snowpack of 1983 which continued to build through the winter and ended up producing serious flooding across the state. At this point, it is far too early to predict the outcome of this year's snowpack, but if it follows a maximum type pattern, the Sevier and southern Utah could end up with snowpacks ranging from 200% to 300% of normal. The smallest snowpack in the state is in central Utah, in the San Pitch valley which is near average at 103%. Precipitation for December was near average for most of the state except southern Utah which had about 150% of normal, bringing seasonal precipitation, (Oct-Dec) to 153%. Soil moisture was substantially recharged from large precipitation events in late fall and early winter. Current soil moisture across the entire state is only about 10% to 15% less than what it was during active snowmelt of last spring. Estimates of soil moisture range from about 50% to 70% of saturation in the upper 24 inches of soil. Low reservoir storage is also a concern with total reservoir storage at 38% of capacity, the same as last year. The area of greatest drought concern is the Bear River with current reservoir storage at of 2% of capacity. Areas that could have high streamflows include the upper Sevier and the Virgin. Streamflow forecasts range from 50% to 150% of average. Surface Water Supply Indices range from 4% on the Bear River, to 82% on the Virgin.

SNOWPACK

January first snowpacks as measured by the NRCS SNOTEL system range from 112% on the Bear to 246% in southwestern Utah. Most areas in northern Utah are comparable to last year, whereas the Uintah Basin and everything south of Salina have substantially greater snowpacks. The Midway Valley SNOTEL site currently has 36 inches of snow water equivalent and its April 1 average peak is only 27 inches. Of some concern are low elevation snowpacks across the state, which are below average. Any outcome is still possible, even in southern Utah, but there is a high probability that the upper Sevier, the Virgin and the Uintah Basin will have at least average snowpacks on April 1 and realistically could be 130% or greater.

PRECIPITATION

Mountain precipitation during December was near average over much of Utah (100%-125%). In southern Utah, precipitation was 150% of average. This brings the seasonal accumulation (Oct-Dec) to 153% of average statewide.


RESERVOIRS

Storage in 41 of Utah's key irrigation reservoirs is at 38% of capacity. This is the same as last year and reflects heavy use of reservoir storage to make up the streamflow deficit during years of drought. Most reservoir operators are utilizing a conservative strategy, storing as much water as possible.


STREAMFLOW

Snowmelt streamflows are expected to be below average to well above average across the state of Utah this year. Forecast streamflows range from 51% on the Bear at Stewart dam to 238% on Coal Creek near Cedar City. Most flows are forecast to be in the 80% to 120% range. Overall water supply conditions are improving.


Mountain Snowpack



Precipitation





Statewide Reservoir Storage



Bear River Basin Jan 1, 2005

Snowpacks on the Bear River Basin are slightly above average at 112% of normal, about 98% of last year. Specific sites range from 60% to 135% of normal. December precipitation was near average at 103%, which brings the seasonal accumulation (Oct-Dec) to 118% of average. Soil moisture levels in runoff producing areas are at 64% of saturation in the upper 2 feet of soil compared to 32% last year. Forecast streamflows range from much below to near average (51%-111%) volumes this spring. Reservoir storage is extremely low at 2% of capacity, the same as last year. The Surface Water Supply Index is at 4% for the Bear River, or 96% of years have had more total water available. Water supply conditions are much below normal due to low reservoir storage..





BEAR RIVER BASIN

Streamflow Forecasts - January 1, 2005

Forecast Point	Forecast	İ	Drier ====	== Future Co		===== Wette:	r ====>>	
	Period	90% (1000AF)	70% (1000AF)	50 (1000AF))% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Bear River nr UT-WY State Line	APR-JUL	86	109	 125	111	141 141	164	113
Bear River ab Reservoir nr Woodruff	APR-JUL	63	100	 125	92	150	189	136
Big Creek nr Randolph	APR-JUL	2.10	3.60	I 4.60	94	 5.60	7.10	4.90
Smiths Fork nr Border	APR-JUL	58	80	I 95	92	110	132	103
Bear River at Stewart Dam	APR-JUL	50	88	 120	51	157	221	234
Little Bear River at Paradise	APR-JUL	23	36	 46	100	I 58	77	46
Logan River nr Logan combined flow	APR-JUL	63	89	 110	87	133	170	126
Blacksmith Fork nr Hyrum	APR-JUL	24	36	I 4 5 	94	I 56 	73	48


		RIVER BASIN	. 6	!	BEAR RIVER BASIN Watershed Snowpack Analysis - January 1, 2005						
	Reservoir Storage (1	000 AF) - End	of Decem	ber =======	ا 	watershed Showpack Analysis - January 1, 200					
Reservoir		Usable Capacity	*** Usable Storage *** Number This Last Watershed of				r as % of				
		1	Year	Year	Avg		Data Sites	Last Yr	Average		
BEAR LAKE		1302.0	0.0	15.7		BEAR RIVER, UPPER (abv	На 6	116	121		
HYRUM		15.3	7.3	6.4	10.2	BEAR RIVER, LOWER (blw	Ha 8	88	106		
PORCUPINE		11.3	6.5	5.0	3.9	LOGAN RIVER	4	94	114		
WOODRUFF NAF	ROWS	57.3	12.0	6.0	23.6	RAFT RIVER	1	81	101		
WOODRUFF CRE	EK	4.0	1.5	1.2	j	BEAR RIVER BASIN	14	98	112		

^{* 90%, 70%, 50%, 30%,} and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
 The value is natural volume - actual volume may be affected by upstream water management.

Weber and Ogden River Basins Jan 1, 2005

Snowpack on the Weber and Ogden Watersheds is above normal at 136% of average, about 100% of last year. Individual sites range from 81% to 188% of average. December precipitation was near average at 109% bringing the seasonal accumulation (Oct-Dec) to 139% of average. Soil moisture levels in runoff producing areas are at 66% of saturation in the upper 2 feet of soil compared to 30% last year. Streamflow forecasts range from 92% to 132% of average. Reservoir storage is at 43% of capacity, about 11% more than last year. The Surface Water Supply Index is at 31% for the Weber River and at 54% for the Ogden River. Overall water supply conditions are below to near normal due to low reservoir storage.



WEBER & OGDEN WATERSHEDS in Utah Streamflow Forecasts - January 1, 2005

APR-JUL

5.30

<<===== Drier ===== Future Conditions Wetter ====>> Forecast Point == Chance Of Exceeding * Forecast Period 90% 70% 30% 10% 30-Yr Avg. 50% (1000AF) (1000AF) (1000AF) (% AVG.) (1000AF) (1000AF) Smith & Morehouse Res inflow APR-JUL 25 31 36 106 41 47 34 Weber River nr Oakley APR-JUL 119 135 110 151 123 Rockport Reservoir inflow APR-JUL 91 123 145 108 167 198 134 127 150 173 137 Weber River nr Coalville APR-JUL 93 110 206 Chalk Creek at Coalville APR-JUL 22 36 45 100 54 68 45 185 179 APR-JUL 117 157 215 255 Echo Reservoir inflow 103 Lost Creek Reservoir inflow APR-JUL 7.1 12.1 16.2 92 21 29 17.6 16.7 37 47 31 East Canyon Reservoir inflow APR-JUL 24 30 97 215 300 355 355 Weber River at Gateway APR-JUL 100 410 495 SF Ogden River nr Huntsville APR-JUL 40 58 70 109 82 100 64 117 133 APR-JUL 84 140 105 163 195 Pineview Reservoir inflow

	WEBER & OGDEN WA Reservoir Storage (1000			ber	 	WEBER & OGDEN WATERSHEDS in Utah Watershed Snowpack Analysis - January 1, 200				
Reservoir		Usable Capacity 	*** Usa This Year	ble Storaç Last Year	Avg	Watershed	Number of Data Sites	This Yea	r as % of Average	
CAUSEY		7.1	3.3	2.1	2.8	OGDEN RIVER	4	86	126	
EAST CANYON		49.5	33.7	23.9	34.9	WEBER RIVER	9	97	142	
ЕСНО		73.9	33.1	35.5	47.9	WEBER & OGDEN WATERS	HEDS 13	94	136	
LOST CREEK		22.5	5.2	1.3	14.1					
PINEVIEW		110.1	65.6	27.6	52.9					
ROCKPORT		60.9	40.7	30.4	36.2 J					
WILLARD BAY		215.0	47.0	42.7	147.7 147.7					

7.10

8.30

132

9.50

11.30

6.30

The average is computed for the 1971-2000 base period.

Wheeler Creek nr Huntsville


^{* 90%, 70%, 50%, 30%,} and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

^{(1) -} The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels. (2) - The value is natural volume - actual volume may be affected by upstream water management.


Utah Lake, Jordan River & Tooele Valley Basins Jan 1, 2005


Snowpacks over these watersheds are at 134% of average, 99% of last year. Individual sites range from 53% to 231% of average. December precipitation was near average at 112%, bringing the seasonal accumulation (Oct-Dec) to 144% of average. Soil moisture levels in runoff producing areas are at 71% of saturation in the upper 2 feet of soil compared to 38% last year. Forecast streamflows range from 74% to 127% of average. Reservoir storage is at 58% of capacity, 3% less than last year. The Surface Water Supply Index is at 19%, or 81% of years would have more total water available. General water supply conditions are below normal due to low reservoir storage.

Provo River Snowpack



Provo River Precipitation





UTAH LAKE, JORDAN RIVER & TOOELE VALLEY Streamflow Forecasts - January 1, 2005

		 <<=====	======= = Drier =		Future Co	nditions ==	Wett	er ====	==>>	
Forecast Point	Forecast Period		70% (1000AF	1	50	(% AVG.)	30% 30% (1000AE	1(r) (10(30-Yr Avg. (1000AF)
Spanish Fork River nr Castilla	APR-JUL	24	48	 	72	94	96		L20	77
Provo River nr Woodland	APR-JUL	69	97		111	108	125	1	L53	103
Provo River nr Hailstone	APR-JUL	70	104	-	121	111	138	1	L72	109
Provo R blw Deer Creek Dam	APR-JUL	64	114	-	141	112	 168	2	220	126
American Fk R nr American Fk	APR-JUL	19.8	31	-	37	116	l 43		54	32
Utah Lake inflow	APR-JUL	156	274	-	350	108	 426	į	545	325
Little Cottonwood Ck nr SLC	APR-JUL	32	40	-	45	113	I I 50		58	40
Big Cottonwood Ck nr SLC	APR-JUL	26	35		40	105	l I 45		54	38
Mill Creek nr SLC	APR-JUL	3.60	5.87		7.00	100	 8.13	10	. 40	7.00
Parley's Creek nr SLC	APR-JUL	5.2	14.8	!	15.3	92	 15.8		25	16.7
Dell Fork nr SLC	APR-JUL	1.77	4.48		6.30	93	 8.12	10	. 90	6.80
Emigration Creek nr SLC	APR-JUL	0.63	2.95		4.40	98	l 5.85	8	.20	4.50
City Creek nr SLC	APR-JUL	4.10	7.12	!	8.90	102	 10.68	13	. 70	8.70
Vernon Creek nr Vernon	APR-JUL	0.61	0.91	!	1.20	81	l 1.58	2	. 37	1.48
Settlement Creek nr Tooele	APR-JUL	0.81	1.16		1.45	74	1.78	2	. 36	1.97
South Willow Creek nr Grantsville	APR-JUL	2.30	3.40		4.10	127	 4.80 	5	. 90	3.23
UTAH LAKE, JORDAN Reservoir Storage (100		ELE VALLEY				UTAH LAKE, Watershed Si	JORDAN RIVE	R & TOO ysis -	DELE VA	
Reservoir	Usable Capacity 	*** Usab This Year	le Storag Last Year		 Water		Nun	ber f Sites		Year as % of
DEER CREEK	149.7	95.2	54.0	102.0	====== PROVO	RIVER & UTA	AH LAKE	7	96	122
GRANTSVILLE	3.3	1.5	0.6	1.6	 PROVO	RIVER		4	100	134
SETTLEMENT CREEK	1.0	0.6	0.5	0.5	 JORDA	N RIVER & G	REAT SALT	6	92	137
STRAWBERRY-ENLARGED	1105.9	735.3	778.9	640.0	 TOOEL 	E VALLEY WA	TERSHEDS	3	97	156

^{* 90%, 70%, 50%, 30%,} and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

0.2

756.5 | UTAH LAKE, JORDAN RIVER & 16

134

The average is computed for the 1971-2000 base period.

UTAH LAKE

VERNON CREEK

870.9 384.6

0.4


0.6


The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
 The value is natural volume - actual volume may be affected by upstream water management.


Uintah Basin and Dagget SCD's Jan 1, 2005

Snowpacks across the Uintah Basin and North Slope areas are much above average and near record levels at 177%, which is 146% of last year. The North Slope ranges from 93% to 276% and the Uintah Basin ranges from 115% to 257% of average. Precipitation during December was above average at 126% bringing the seasonal accumulation (Oct-Dec) to 170% of average. Soil moisture estimates in runoff producing areas are at 55% of saturation in the upper 2 feet of soil compared to 30% last year. Reservoir storage is at 67% of capacity, 2% less than last year. The Surface Water Supply Index for the western area is 70% and for the eastern area it is 67% indicating slightly above normal conditions basin wide. Streamflow forecasts range between 108% and 157% of average. Springtime runoff conditions are above normal. There is a 97% probability of getting at least average snowpack by April 1 and a high probability of snowpacks greater than 130% of normal. Preparation for high flows might be considered.

Uintahs Snowpack







UINTAH BASIN & DAGGET SCD'S Streamflow Forecasts - January 1, 2005

		<<===== 	Drier ====	== Future Co	nditions ==	===== Wetter	====>>	:======== : (
Forecast Point	Forecast Period	====== 90% (1000AF)	70% (1000AF)	50	% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Blacks Fork nr Robertson	APR-JUL	65	83	96	101	109	127	95
EF of Smiths Fork nr Robertson	APR-JUL	22	27	30	97	34	41	31
Flaming Gorge Reservoir Inflow	APR-JUL	675	930	1100	92	1270	1530	1190
BIG BRUSH CK abv Red Fleet Resv	APR-JUL	18.0	23	26	124	 29	34	21
Ashley Creek nr Vernal	APR-JUL	52	66	75	144	 84	98	52
WF DUCHESNE RIVER nr Hanna	APR-JUL	16.0	23	28	117	 34	43	24
DUCHESNE R nr Tabiona	APR-JUL	79	98	110	105	122	141	105
UPPER STILLWATER RESV inflow	APR-JUL	69	87	100	122	113	131	82
ROCK CK nr Mountain Home	APR-JUL	82	98	110	124	122	138	89
DUCHESNE R abv Knight Diversion	APR-JUL	145	187	215	114	245	285	188
STRAWBERRY RES nr Soldier Springs	APR-JUL	28	48	65	110	 84	117	59
CURRANT CREEK RESV Inflow	APR-JUL	17.8	23	27	108	31	36	25
STARVATION RESERVOIR inflow	APR-JUL	69	111	140	116	1 169	211	121
Lake Fork River abv Moon Lake	APR-JUL	68	81	90	132	 99	112	68
Yellowstone River nr Altonah	APR-JUL	62	78	88	142	 98	114	62
DUCHESNE R at Myton	APR-JUL	230	310	360	139	 410	490	260
Whiterocks River nr Whiterocks	APR-JUL	53	70	82	146	 94	111	56
DUCHESNE R nr Randlett	APR-JUL	260	410	510	157	 610	760	325

	UINTAH BASIN & DAGGET S Storage (1000 AF) - End	d of Dece			Watershed Snowpack A	UINTAH BASIN & DAGGET SCD'S Watershed Snowpack Analysis - January 1,				
Reservoir	Usable Capacity	*** Us	able Stora Last Year		Watershed	Number of ta Sites	This Yea	r as % of 		
FLAMING GORGE	3749.0	2741.0	2605.0	3027.0	UPPER GREEN RIVER in UTAH	6	143	159		
MOON LAKE	49.5	17.2	13.5	26.1	ASHLEY CREEK	2	168	228		
RED FLEET	25.7	15.4	12.9	17.5	BLACK'S FORK RIVER	2	115	105		
STEINAKER	33.4	14.6	9.5	20.0	SHEEP CREEK	1	157	152		
STARVATION	165.3	123.4	123.1	128.6	DUCHESNE RIVER	11	147	184		
STRAWBERRY-ENLARGED	1105.9	735.3	778.9	640.0	LAKE FORK-YELLOWSTONE CRE	4	178	198		
					STRAWBERRY RIVER	4	103	144		
					UINTAH-WHITEROCKS RIVERS	2	192	250		
					UINTAH BASIN & DAGGET SCD	17	146	177		


^{* 90%, 70%, 50%, 30%,} and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

^{(1) -} The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.(2) - The value is natural volume - actual volume may be affected by upstream water management.


Carbon, Emery, Wayne, Grand and San Juan Co. Jan 1, 2005


Snowpacks in this region are much above normal at 140% of average, about 123% of last year. Individual sites range from 112% to 314% of average. Precipitation during December was near average at 101%, bringing the seasonal accumulation (Oct-Dec) to 150% of normal. Soil moisture estimates in runoff producing areas are at 60% of saturation in the upper 2 feet of soil compared to 30% last year. Forecast streamflows range from 83% to 108% of average. Reservoir storage is at 35% of capacity, down 2% from last year. Surface Water Supply Indices for the area are: Price 25%, (much below normal) San Rafael area 56% (near average) and Moab 50% (near average). General runoff and water supply conditions are below to near normal.

Southeast Utah Snowpack



Southeast Utah Precipitation





CARBON, EMERY, WAYNE, GRAND, & SAN JUAN Co.

	1	<<=====	= Drier ==	==== :	Future Co	nditions ==	==== Wetter	====>>	
Forecast Point		90% (1000AF)	70% (1000AF)	 	50 (1000AF)	% (% AVG.)	30% (1000AF)	10% (1000AF)	
coseberry Creek nr Scofield	APR-JUL	5.8	9.2	== ===: 	11.5	97	13.8	17.2	11.9
cofield Reservoir inflow	APR-JUL	27	37	l I	44	96	51	61	46
Thite River blw Tabbyune Creek	APR-JUL	8.6	14.2	l I	18.8	108	24	33	17.4
reen River at Green River, UT	APR-JUL	1790	2570	l I	3100	98	3630	4410	3170
lectric Lake inflow	APR-JUL	6.6	10.9	l I	14.6	93	19.1	27	15.7
UNTINGTON CK nr Huntington	APR-JUL	24	37	l I	46	92	55	68	50
OE'S VALLEY RESV Inflow	APR-JUL	27	43	l I	54	93	65	81	58
erron Creek nr Ferron	APR-JUL	22	30		37	95	44	56	39
colorado River nr Cisco	APR-JUL	2410	3540	l l	4300	93	5060	6190	4650
Mill Creek at Sheley Tunnel nr Moab	APR-JUL	2.20	4.50	l I	6.00	120	7.50	9.80	5.00
even Mile Creek nr Fish Lake	APR-JUL	3.20	5.50	l I	7.10	101	8.70	11.00	7.00
Nuddy Creek nr Emery	APR-JUL	7.7	14.3	l l	18.7	94	23	30	19.9
North Ck ab R.S. nr Monticello	MAR-JUL	0.22	0.70		1.40	144	2.00	3.70	0.97
outh Ck ab Lloyd's Res nr Monticell	MAR-JUL	0.84	1.60		2.00	146	3.20	4.90	1.37
ecapture Ck bl Johnson Ck nr Blandi	MAR-JUL	2.10	5.10	l I	7.50	149	13.10	20.00	5.05
an Juan River nr Bluff	APR-JUL	930	1250	 	1460	119 	1670	1990	1230
CARBON, EMERY, WAYNE, Reservoir Storage (1000	GRAND, & S AF) - End	AN JUAN Co	o. er		l C	ARBON, EMERY Watershed Sno	, WAYNE, GRAN owpack Analys	ID, & SAN JU	JAN Co. cy 1, 2005
deservoir	Usable Capacity		le Storage Last		 Water		Numbe of	r This	Year as % o
	- 1	Year	Year	Avg	İ	snea	Data Si	tes Last	
UNTINGTON NORTH	4.2	2.2	3.1	2.4	1	RIVER	3	102	123
TOE'S VALLEY	61.6	35.3	32.5	41.0	I SAN R	AFAEL RIVER	3	106	118
EN'S LAKE	2.3	0.3	0.5	1.0	MUDDY	CREEK	1	90	113
IILL SITE	16.7	8.5	6.9	75.0	 FREMO	NT RIVER	3	217	217
COFIELD	65.8	7.0	12.8	32.7	 LASAL	MOUNTAINS	1	93	119
					 BLUE	MOUNTAINS	1	173	161
					I		1	93	134

^{* 90%, 70%, 50%, 30%,} and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

CARBON, EMERY, WAYNE, GRA 13 123 140


⁽¹⁾ - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels. (2) - The value is natural volume - actual volume may be affected by upstream water management.


Sevier and Beaver River Basins Jan 1, 2005


Snowpacks on the Sevier River Basin are much above normal at 172% of average, about 143% of last year. The lower Sevier area is near average at 103%. Individual sites range from 79% to 374% of average. Precipitation during December was near average at 102% of normal, bringing the seasonal accumulation (Oct-Dec) to 182% of average. Soil moisture estimates in runoff producing areas are at 62% of saturation (Sevier) and 21% (Beaver) in the upper 2 feet of soil compared to 31% last year. Streamflow forecasts range from 96% to 140% of average. Reservoir storage is at 18% of capacity, 3% more than last year. Surface Water Supply Indices are: Upper Sevier 50%, Lower Sevier 60% and Beaver 54%. Water supply conditions are near average due to high snowpack and soil moisture. The Sevier has an 85% probability of at least average snowpacks on April 1 and significant probability of 130% or more. On the upper Sevier, preparation for high flows might be appropriate.

Sevier River Precipitation

Sevier River Snowpack







SEVIER & BEAVER RIVER BASINS Streamflow Forecasts - January 1, 2005

	!	<<=====	Drier ===	1	Future Co	onditions ==	===== We	tter ===	==>>	
Forecast Point	Forecast	======		=== Cha	ance Of E	Exceeding * =			====	
	Period	90%	70%	1	50)% [30%			30-Yr Avg.
	l	(1000AF)	(1000AF)	_ !	(1000AF)	(% AVG.)	(1000	AF) (10	00AF)	(1000AF)
Sevier River at Hatch	APR-JUL	25	61	 !	77	140	9	4	101	55
Sevier River nr Kingston	APR-JUL	51	101	-	121	136 I	14	1	148	89
EF Sevier R nr Kingston	APR-JUL	19.0	46	-	56	147	6	7	71	38
Sevier R blw Piute Dam	APR-JUL	76	146	-	177	141	20	8	215	126
Clear Creek nr Sevier	APR-JUL	9.9	26		31	141	3	6	38	22
Salina Creek at Salina	APR-JUL	5.2	18.2	-	27	137	3	6	49	19.7
Sevier R nr Gunnison	APR-JUL	78	281	!	365	130	44	9	510	280
Chicken Creek nr Levan	APR-JUL	1.42	2.88	!	4.30	96 I	6.1	.3 9	. 65	4.50
Oak Creek nr Oak City	APR-JUL	0.96	1.42	!	1.78	107	2.1	.8 2	2.85	1.66
Beaver River nr Beaver	APR-JUL	24	31	-	37	137	4	3	54	27
Minersville Reservoir inflow	APR-JUL	8.9	15.5		21	127 	2	7	38	16.6
SEVIER & BEAN	ER RIVER BAS	INS			 I	SEVIER	& BEAVER	RIVER E	BASINS	
Reservoir Storage (100	00 AF) - End	of Decembe	er		I	Watershed Sn	owpack An	alysis -	January	L, 2005
	Usable	*** Usab]	Le Storage	***	 I		N	umber	This Yea	ar as % of
Reservoir	Capacity 	This Year	Last Year	Avg	Water 	rshed	Dat	of a Sites	====== Last Yr	Average
GUNNISON	20.3	1.1	0.0	10.9	====== UPPEF	SEVIER RIVE		8	201	251
MINERSVILLE (RkyFd)	23.3	5.0	3.6	12.7	 EAST	FORK SEVIER	RIVER	3	210	243
OTTER CREEK	52.5	10.7	13.2	32.8	 SOUTE	H FORK SEVIER	RIVER	5	195	255
PIUTE	71.8	18.0	16.4	42.1	 LOWEF	R SEVIER RIVE	R (inclu	6	86	103
SEVIER BRIDGE	236.0	37.9	28.0 1	L48.9	 BEAVE	ER RIVER		2	145	155
PANGUITCH LAKE	22.3	6.5	3.8 1	108.0	 SEVIE	ER & BEAVER R	RIVER BAS	16	147	172
					I					


^{* 90%, 70%, 50%, 30%,} and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
 The value is natural volume - actual volume may be affected by upstream water management.


E. Garfield, Kane, Washington, & Iron co. Jan 1, 2005


Snowpacks in this region are much above normal at 246% of average, about 229% of last year. Individual sites range from 133% to 374% of average. Precipitation was much above normal during December at 150% of average, bringing the seasonal accumulation (Oct-Dec) to 265% of normal. Soil moisture estimates in runoff producing areas are at 70% of saturation in the upper 2 feet of soil compared to 30% last year. Forecast streamflows range from 172% to 238% of average. Reservoir storage is at 59% of capacity, 18% more than last year. The Surface Water Supply Index is at 82%, indicating much above normal water availability. While this is only January, concerns over the potential for high flows this spring are increasing. This area has an 88% probability of at least average snowpacks on April 1 and significant potential of snowpacks of 150%.

Southwest Utah Snowpack



Southwest Utah Precipitation





E. GARFIELD, KANE, WASHINGTON, & IRON Co. Streamflow Forecasts - January 1, 2005

		Streamil	.ow Forecast							
		<<==== 	== Drier ==	====	Future Co	nditions ==	===== V	letter	====>>	I I
Forecast Point	Forecast Period	===== 90% (1000AE	70% (1000AF)	I I	50 (1000AF)	* xceeding * =	30 (100)%)OAF)	10% (1000AF)	 30-Yr Avg. (1000AF)
Lake Powell inflow	APR-JUL	4320	6390	==!===: !	7800	98		10	11280	7930
Virgin River nr Virgin	APR-JUL	58	85	!	110	172	1	.34	162	64
Virgin River nr Hurricane	APR-JUL	43	106	-	121	175	1	.36	199	69
Santa Clara River nr Pine Valley	APR-JUL	3.40	7.42	-	10.00	182	12.	96	16.60	5.50
Coal Creek nr Cedar City	APR-JUL	29	39		46	238 		54	67	19.3
E. GARFIELD, KANE, Reservoir Storage (10	,			=====		E. GARFIELD, Watershed Sn				
Reservoir	Usable Capacity	This	ble Storage Last		======= Water	shed		Number of		Year as % of
	1	Year	Year	Avg	I		Da	ta Sit	es Last	Yr Average
GUNLOCK	10.4	Year ======== 10.4	Year 	Avg ===== 5.7		N RIVER				3 -
	10.4		3.8			N RIVER				
GUNLOCK		10.4	3.8	5.7	VIRGI PAROW	N RIVER		5	210	263
GUNLOCK LAKE POWELL	24322.0	10.4	3.8	5.7	VIRGI PAROW ENTER	IN RIVER		5	210 265	263 333

^{* 90%, 70%, 50%, 30%,} and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

E. GARFIELD, KANE, WASHIN 9

227

246

The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
 The value is natural volume - actual volume may be affected by upstream water management.

UTAH SURFACE	WATER	SUPPLY	INDEX
Snow Surveys	NRCS	USDA	
Basin or Region	SWSI/%	Percentile	Years with
1-Jan-05			Similar SWSI
Bear River	-3.8	4%	04,03,93
Ogden River	0.3	54%	89,93,96,95
Weber River	-1.6	31%	87,00,89,94
Provo	-2.6	19%	56,03,55,59
West Uintah Basin	1.6	70%	02,96,86,01
East Uintah Basin	1.4	67%	93,85,99,84
Price River	-2.1	25%	59,02,03,89
San Rafael	0.5	56%	00,74,82,98
Moab	0.0	50%	82,91,94,97
Upper Sevier River	0.0	50%	76,75,01,74
Lower Sevier River	0.8	60%	00,75,81,70
Beaver River	0.4	54%	96,78,74,81
Virgin River	2.6	82%	92,88,98,95
0			0)4/01 0 1 4 + - 4
Snow Surveys 245 N Jimmy Doolittle Rd			SWSI Scale: -4 to 4 Percentile: 0 - 100%
Salt Lake City, UT			rercentile: U - 100%
(801) 524-5213			
(001) 024-0210			

What is a Surface Water Supply Index?

The Surface Water Supply Index (SWSI) is a predictive indicator of total surface water availability within a watershed for the spring and summer water use seasons. The index is calculated by combining pre-runoff reservoir storage (carryover) with forecasts of spring and summer streamflow which are based on current snowpack and other hydrologic variables. SWSI values are scaled from +4.1 (abundant supply) to -4.1 (extremely dry) with a value of zero (0) indicating media water supply as compared to historical analysis. SWSI's are calculated in this fashion to be consistent with other hydroclimatic indicators such as the Palmer Drought Index and the Precipitation index.

Utah Snow Surveys has also chosen to display the SWSI as a PERCENT CHANCE OF NON-EXCEEDANCE. While this is a very cumbersome name, it has the simplest application. It can be best thought of as a simple scale of 1 to 99 with 1 being the drought of record (driest possible conditions) and 99 being the flood of record (wettest possible conditions) and a value of 50 representing average conditions. This rating scale is a percentile rating as well, for example a SWSI of 75% means that this years water supply if greater than 75% of all historical events and that only 25% of the time has it been exceeded. Conversely a SWSI of 10% means that 90% of historical events have been greater than this one and that only 10% have had less total water supply. This scale is far more intuitive for most people and is totally comparable between basins: a SWSI of 50% means the same relative ranking on watershed A as it does on watershed B, which may not be strictly true of the +4 to -4 scale.

In general terms, the scale is divided into 3 parts: LOW - AVERAGE - HIGH. 1 to 33 means below normal water supply, 34 to 66 means average conditions and 67 to 99 means above average water supply amounts.

Another benefit to the SWSI is in the calculation table or the actual ranking of each individual year. The current SWSI can be directly compared to the years it most closely resembles. In other words, is the current year similar to the familiar floods of 1983 or the drought of 1977. Managers can then refer to records from those particular years in determining strategies for dealing with the current years water supply. Also evident is whether the streamflow component or the reservoir component is the predominant driving force at any given time.

SWSI's can be an excellent water management tool in determining over all risk and management strategies. It gives the water user and manager more information than simply streamflow or reservoir level alone.

SNOW COURSE DATA

JANUARY 2005

SNOW COURSE	ELEV.	DATE	SNOW	WATER	LAST	AVERAGE
			DEPTH		YEAR	71-00
AGUA CANYON SNOTEL	8900	1/01	 35	7.4	3.0	2.9
ALTA CENTRAL	8800	12/28	63	20.2	20.0	16.5
BEAVER DAMS SNOTEL	8000	1/01	18	3.4	4.3	4.3
BEAVER DIVIDE SNOTEL	8280	1/01	34	4.9	4.8	4.7
BEN LOMOND PK SNOTEL	8000	1/01	-	22.4	23.6	14.5
BEN LOMOND TR SNOTEL	6000	1/01	36	8.6	16.3	8.5
BEVAN'S CABIN	6450				-	4.2
BIG FLAT SNOTEL	10290	1/01	56	12.6	7.5	7.6
BIRCH CROSSING	8100	- 10-				2.8
BLACK FLAT-U.M. CK S		1/01	28	5.8	5.4	3.8
BLACK'S FORK GS-EF	9340				-	3.3
BLACK'S FORK JUNCTN BOX CREEK SNOTEL	8930	1 /01	43	0.4		3.7 5.3
BRIAN HEAD	9800 10000	1/01	43	9.4	7.3	8.2
BRIGHTON SNOTEL	8750	1/01	70	13.7	11.2	10.9
BRIGHTON CABIN	8700	12/28	47	13.7	13.2	11.5
BROWN DUCK SNOTEL	10600	1/01	86	17.0	8.1	7.7
BRYCE CANYON	8000	1,01	00	27.0	2.8	2.1
BUCK FLAT SNOTEL	9800	1/01	36	8.3	8.6	7.2
BUCK PASTURE	9700	1,01	50	0.5	-	-
BUCKBOARD FLAT	9000				_	5.4
BUG LAKE SNOTEL	7950	1/01	47	8.8	9.2	8.3
BURT'S-MILLER RANCH		-,			_	2.2
CAMP JACKSON SNOTEL		1/01	27	9.0	5.2	5.6
CASCADE MOUNTAIN SNO	7770	1/01	29	8.9	12.0	_
CASTLE VALLEY SNOTEL	9580	1/01	59	12.6	6.4	4.9
CHALK CK #1 SNOTEL	9100	1/01	62	12.9	9.7	10.1
CHALK CK #2 SNOTEL	8200	1/01	42	7.3	6.9	6.7
CHALK CREEK #3	7500				-	3.5
CHEPETA SNOTEL	10300	1/01	82	15.4	6.8	6.0
CLAYTON SPRINGS SNTL	10000	1/01	51	11.7	5.4	_
CLEAR CK RIDG #1 SNT	9200	1/01	48	9.6	9.3	7.7
CLEAR CK RIDG #2 SNT		1/01	36	4.6	7.4	6.0
CORRAL	8200				-	_
CURRANT CREEK SNOTEL		1/01	35	5.8	6.4	4.2
DANIELS-STRAWBERRY S		1/01	42	9.0	9.4	6.5
DILL'S CAMP SNOTEL	9200	1/01	38	6.2	6.9	5.5
DONKEY RESERVOIR SNO		1/01	36	6.9	2.4	4.0
DRY BREAD POND SNTL	8350	1/01	50	9.2	11.6	9.1
DRY FORK SNOTEL	7160	1/01	18	4.0	8.8	6.9
EAST WILLOW CREEK SN FARMINGTON U. SNOTEL		1/01 1/01	26 69	3.9 24.5	4.2 23.8	2.9 13.0
FARMINGTON U. SNOTEL FARMINGTON LOWER SC	6950	1/01	09	24.5	23.6	10.4
FARMINGTON L. SNOTEL		1/01	39	11.6	19.4	-
FARNSWORTH LK SNOTEL	9600	1/01	41	9.9	8.9	8.0
FISH LAKE	8700	1,01	71	3.3	-	2.9
FIVE POINTS LAKE SNO		1/01	82	14.9	8.6	7.0
G.B.R.C. HEADQUARTER		-,			_	-
	10000				_	9.7
GARDEN CITY SUMMIT	7600				_	6.5
GARDNER PEAK SNOTEL		1/01	37	8.9	-	_
GEORGE CREEK	8840				-	-
GOOSEBERRY R.S.	8400				-	5.1
GOOSEBERRY R.S. SNTL	7900	1/01	14	3.3	5.3	3.6
GUTZ PEAK SNOTEL	6820	1/01	30	8.9	-	_
HARDSCRABBLE SNOTEL		1/01	40	9.8	14.0	6.5
HARRIS FLAT SNOTEL		1/01	21	5.2	3.3	2.5
HAYDEN FORK SNOTEL	9100	1/01	46	8.5	5.8	6.3
HENRY'S FORK	10000		_			
HEWINTA SNOTEL	9500		21	3.8	4.1	
HICKERSON PARK SNTL			21	4.4	2.8	2.9
HIDDEN SPRINGS		12/28	5	2.0	6.8	.2
HOBBLE CREEK SUMMIT		- /0-	0.0	2 2	-	6.1
HOLE-IN-ROCK SNOTEL			22	3.9	3.2	
HORSE RIDGE SNOTEL		1/01	56	12.3	10.3	9.3
HUNTINGTON-HORSESHOE		1 /01	20	۵.0	- 6 1	9.7
INDIAN CANYON SNOTEL SNOW COURSE			38 SMOM	9.0 พลฑะอ	6.1	4.4
SHOW COURSE	ELEV.	DATE	SNOW	WATER	LAST	AVERAGE

			DEPTH	CONTENT	YEAR	71-00
JOHNSON VALLEY	8850				-	2.7
JONES CORRAL G.S.	9720				-	-
KILFOIL CREEK	7300	10/00		0 =	-	5.5
KILLYON CANYON KIMBERLY MINE SNOTEL	6300 9300	12/29 1/01	12 36	2.7 7.2	9.4 7.4	5.1 6.0
KING'S CABIN SNOTEL	8730	1/01	40	9.4	7.2	5.0
KLONDIKE NARROWS	7400	,			-	7.5
KOLOB SNOTEL	9250	1/01	68	16.6	8.8	6.9
LAKEFORK #1 SNOTEL	10100	1/01	66	11.8	6.3	5.6
LAKEFORK BASIN SNTL LAKEFORK MOUNTAIN #3	10900 8400	1/01	76	12.8	8.8	8.2 2.8
LAMBS CANYON	7400	12/29	32	7.4	_	7.4
LASAL MOUNTAIN LOWER	8800				-	3.8
LASAL MOUNTAIN SNTL	9850	1/01	21	5.6	6.0	4.7
LIGHTNING RIDGE SNTL	8220	1/01	44	9.6	-	-
LILY LAKE SNOTEL LITTLE BEAR LOWER	9050 6000	1/01	35	6.2	5.6 -	5.5 4.3
LITTLE BEAR SNOTEL	6550	1/01	24	5.3	8.5	5.2
LITTLE GRASSY SNOTEL	6100	1/01	-	2.8	0.2	2.1
LONG FLAT SNOTEL	8000	1/01	-	4.5	2.5	2.8
LONG VALLEY JCT. SNT	7500	1/01	14	2.9	3.4	1.8
LOOKOUT PEAK SNOTEL LOST CREEK RESERVOIR	8200 6130	1/01	58	15.2	18.0	9.9 2.0
LOUIS MEADOW SNOTEL	6700	1/01	30	6.9	14.7	-
MAMMOTH-COTTONWD SNT	8800	1/01	31	8.3	9.0	7.6
MERCHANT VALLEY SNTL	8750	1/01	34	7.5	6.4	5.4
MIDDLE CANYON	7000	1 (01			-	5.9
MIDWAY VALLEY SNOTEL MILL CREEK	9800 6950	1/01 12/29	90 27	33.7 6.8	11.1 -	9.0 8.3
MILL-D NORTH SNOTEL	8960	1/01	52	16.0	13.7	10.3
MILL-D SOUTH FORK	7400	12/28	31	8.0	11.7	8.6
MINING FORK SNOTEL	8000	1/01	43	12.7	11.1	5.5
MONTE CRISTO SNOTEL	8960	1/01	62	14.1	11.7	11.0
MOSBY MTN. SNOTEL MT.BALDY R.S.	9500 9500	1/01	56	12.3	7.6	5.1 9.9
MUD CREEK #2	8600				<u>-</u>	5.3
OAK CREEK	7760				=	-
PANGUITCH LAKE R.S.	8200				-	-
PARLEY'S CANYON SNTL	7500	1/01	34	5.8	9.7	7.2
PARRISH CREEK SNOTEL	7740	1/01	46	12.7	17.9	- 7.2
PAYSON R.S. SNOTEL PICKLE KEG SNOTEL	8050 9600	1/01 1/01	40 26	7.4 5.4	8.4 7.4	6.2
PINE CREEK SNOTEL	8800	1/01	-	9.4	11.0	8.8
RED PINE RIDGE SNTL	9200	1/01	39	7.5	7.3	6.7
REDDEN MINE LOWER	8500				-	6.7
REES'S FLAT ROCK CREEK SNOTEL	7300 7900	1 /01		4.8	4.8	5.6 3.7
ROCKY BN-SETTLEMT SN		1/01 1/01	- 41	12.6	11.6	10.0
SEELEY CREEK SNOTEL		1/01	35	8.1	6.7	6.4
SMITH MOREHOUSE SNTL	7600	1/01	34	6.4	5.6	5.7
SNOWBIRD SNOTEL	9700	1/01	90	25.1	18.3	13.2
SPIRIT LAKE	10300				-	5.5
SQUAW SPRINGS STEEL CREEK PARK SNO	9300 10100	1/01	38	7.5	- 5.7	3.2 6.7
STILLWATER CAMP	8550	-, V-	50		-	3.9
STRAWBERRY DIVIDE SN		1/01	48	8.5	9.5	7.4
SUSC RANCH	8200				-	2.8
TALL POLES	8800	1 /01	42	0.4	- 0 1	5.3
TEMPLE FORK SNOTEL THAYNES CANYON SNTL	7410 9200	1/01 1/01	43 78	8.4 18.1	8.4 11.0	- 9.0
THISTLE FLAT	8500	-, V-	. 0	-3.1	-	-
TIMBERLINE	9100				=	_
TIMPANOGOS DIVIDE SN		1/01	62	13.7	13.2	9.2
TONY GROVE LK SNOTEL		1/01	71	18.0	17.1	14.3
TONY GROVE R.S. TRIAL LAKE	6250 9960				_	5.0 9.8
TRIAL LAKE SNOTEL	9960	1/01	68	13.8	10.7	10.5
TROUT CREEK SNOTEL	9400	1/01	48	11.6	5.3	4.2
UPPER JOES VALLEY	8900				_	4.1
VERNON CREEK SNOTEL		1/01	30	5.1	8.7	4.0
VIPONT WEBSTER FLAT SNOTEL	7670 9200	1/01	39	10.6	- 6.3	- 6.0
WHITE RIVER #1 SNTL		1/01	39 36	7.3	6.4	5.2
WHITE RIVER #3	7400				-	3.5
WIDTSOE #3 SNOTEL	9500	1/01	48	13.8	4.4	4.4
WRIGLEY CREEK	9000				=	4.3
YANKEE RESERVOIR	8700				-	3.7



Issued by

Bruce I. Knight
Chief
Natural Resources Conservation Service
U.S. Department of Agriculture

Prepared by

Snow Survey Staff
Randall Julander, Supervisor
Ray Wilson, Hydrologist
Timothy Bardsley, Hydrologist
Mike Bricco, Hydrologist
Bob Nault, Electronics Technician

Released by

Sylvia Gillen State Conservationist Natural Resources Conservation Service Salt Lake City, Utah

YOU MAY OBTAIN THIS PRODUCT AS WELL AS CURENT SNOW, PRECIPITATION, TEMPERATURE AND SOIL MOISTURE, RESERVOIR, SURFACE WATER SUPPLY INDEX, AND OTHER DATA BY VISITING OUR WEB SITE @:

http://www.ut.nrcs.usda.gov/snow/

Snow Survey, NRCS, USDA 245 North Jimmy Doolittle Road Salt Lake City, UT 84116 (801) 524-5213



Utah Water Supply Outlook Report

Natural Resources Conservation Service Salt Lake City, UT

